



You are scheduled for: colonoscopy upper endoscopy flexible sigmoidoscopy on:

_____ / _____ : _____ AM / PM _____ : _____ AM/PM with:
Day Date Arrival time Procedure Time

Dr. Banerjee Dr. Root Dr. Singh Dr. Thyssen Dr. Weinstock

Your procedure is scheduled at the **Advanced Endoscopy Center** 11525 Olde Cabin Road in Creve Coeur.

- **Notify our office immediately if you are on dialysis, on home oxygen, have a ventricular defibrillator, weigh over 350 pounds or have a BMI over 48. Your procedure will need to be rescheduled to a hospital.**
- Please do **NOT** arrive any earlier than your scheduled **arrival** time. Procedure center doors do not open until 6:00am.
- One week prior to your procedure a nurse from the **Advanced Endoscopy Center** will contact you to pre-register. At that time please have a list of your medications and medical history available for this call. It is very important that you preregister with a nurse prior to the day of your procedure. If you have **NOT** received a call within 3 days prior to your procedure you will need to call **(314) 400-9999 option 2**, Monday through Friday between 1PM-4PM.
- It is facility policy you may **NOT** drive yourself, take a cab, Uber or Lyft home from the Procedure Center. A responsible party will need to stay with you during your entire procedure. You may use a medical transportation company.
- Please bring your **insurance cards, picture ID, co-pay/deductibles.**
- Please call your insurance company to ensure that we participate in your plan.
- If your insurance company requires a referral form or number, it is important that you have obtained this in advance from your primary doctor's office prior to this appointment.
- You will not be able to return to work until the day after the procedure.

Our aim is to provide personalized, caring treatment to people with gastrointestinal conditions and diseases using the most advanced and comprehensive therapies available.

For more information you can visit www.GIdoctor.net or **for questions please call 314-997-0554 option 4.**

Due to ALLERGY and SAFETY reasons patients and their drivers please DO NOT wear perfume, cologne, aftershave or scented body lotions or sprays of any kind.



Directions:

From Interstate I-270:

- Take Olive Blvd. East 0.5 miles to Old Ballas Rd.
- Turn Right on Old Ballas Rd. and drive 0.3 miles to 3-way stop sign
- Turn Left onto Olde Cabin Rd. and turn Left again into the first driveway

From Ballas Rd. (Also called North Ballas Rd.):

- Turn East on Old Ballas Rd. and drive 0.3 miles to a 3-way stop sign
- Drive straight onto Olde Cabin Rd. and turn left into the first driveway

Please note:

- Old Ballas is an "L" shaped street
- When driving on Olive Blvd., Old Ballas is one stoplight east of New Ballas N.
- When driving on New Ballas N., Old Ballas is three streetlights south of Olive Blvd.

Specialists in Gastroenterology
 11525 Olde Cabin Road
 Creve Coeur, Missouri 63141
 Phone: (314) 997-0554
 Fax: (314) 997-5086

Advanced Endoscopy Center
 11525 Olde Cabin Road
 Creve Coeur, Missouri 63141
 Phone: (314) 400-9999
 Fax: (314) 400-9990

Colonoscopy Preparation Using Suflave 2 Day PM/AM Split Dosing Regimen

Cancellation Policy: We require 7-day notice, or you may be charged \$350 cancellation fee. Patients who no-show or late cancel repeatedly, may be denied future appointments. This is not only a courtesy to your physician but provides an opportunity for another patient in need of medical care. Special unavoidable circumstances may have an impact on fees being waived but only with management approval.

If these instructions are not followed it is at the discretion of the physician and your procedure may be cancelled.

• **SEVEN DAYS BEFORE THE PROCEDURE**

Medication restrictions – Please call the prescribing physician

1. *Diabetics taking insulin:* will need dose adjustment the day before the exam. Please ask for advice from the physician that prescribes your insulin. For diabetics only taking diabetic pills simply stop the pills the day before the procedure (see below).
2. *Blood thinners (antithrombotic agents) including: Coumadin (Warfarin), Plavix, Pradaxa, Xarelto, Eliquis, Brilinta and Effient:* These can increase the risk of bleeding during and after endoscopic procedures to a varying degree. **If you take a blood thinner, review our recommendations below for holding these medications prior to your procedure and confirm with your prescribing provider:**

Effient - 7 days; Coumadin (Warfarin), Plavix and Brilinta - 5 days; Pradaxa, Eliquis and Xarelto - 2 days;

**** IF these medications are unable to be stopped, you need to call our office ASAP****

3. Aspirin does **NOT** need to be stopped.

Items to Purchase

1. Fill the prescription for *Suflave*. *Suflave* contains advantame similar to aspartame; if this is a problem, call us before filling the prescription.
2. Fill the prescription for *Ondansetron* (anti-nausea medication).
3. Purchase four (4) Dulcolax tablets "Gentle Overnight Laxative" (over the counter).
4. Any over-the-counter hemorrhoid ointment for anal discomfort (A & D ointment or any diaper cream).
5. Optional water flavor enhancer of your choosing to flavor prep if not using provided lemon lime packet.

Diet restrictions – 7 days prior

No seeds, nuts, popcorn or peels from fruits and vegetables. Examples but not limited to: no peels and seeds from apple, potatoes, cucumbers, zucchini, squash, tomatoes, peppers, corn, no berries (blueberries, raspberries, strawberries), no grapes or bananas etc.

• **TWO DAYS BEFORE THE PROCEDURE**

1. Take two Dulcolax tablets at 6:00pm.

• **ONE DAY BEFORE THE PROCEDURE**

STARTING WHEN YOU WAKE UP, CLEAR LIQUID DIET ONLY FOR THE ENTIRE DAY

1. Drink as much as you can to prevent dehydration. You may have bouillon, broth, black coffee (sugar and artificial sweeteners are okay, no creamer of any kind), tea (honey is okay), juice (apple, grape, cranberry), Gatorade, Powerade, lemonade, soda (regular or diet are okay), water, Popsicles and Jell-O (any flavor or color). Taking red Jell-O and red Gatorade may turn your bowel movements red, do not be alarmed and should be limited. If you have had dehydration before, drink Pedialyte.
2. You may have clear candy (such as Lifesavers, Jolly Ranchers and gummy bears).
3. **DO NOT TAKE** the following diabetic medications: Glucotrol (glipizide), Glyburide (micronase, diabetia), Januvia (sitagliptin), Prandin (repaglinide), Starlix (nateglinide), Symlin (pramlinide).

TIMING OF PREPARATION

Please follow these instructions and NOT the Suflave box instructions

• EVENING BEFORE THE PROCEDURE

This is divided into two sessions. You must drink the entire Suflave to ensure your colon will be cleaned out. A poor prep can mean missed lesions (flat polyps) and cancellation and rescheduling of your procedure.

Between 4p.m. and 7p.m. (first dose)

1. Take the Ondansetron (anti-nausea) 30 minutes prior to taking Dose 1 of the Suflave (bowel prep).
2. Fill first provided bottle with either provided flavor packet (lemon lime flavor) or any water enhancers of your choice like Crystal Light, MiO or Water Drops. Add lukewarm water to the top line of the container (at least 32 ounces), mix with spoon or shake until completely dissolved. You can mix a few hours prior to drinking and refrigerate to improve taste.
3. Drink **ALL** the solution, 8 ounces every 15 mins. This can be done gradually to decrease the risk of nausea.
4. Drink an additional 16 oz or more of a clear liquid of your choice.
5. Take two (2) Dulcolax tablets 1 hour after finishing the first dose of the prep
6. You can continue the clear liquid diet after you finish the first dose of the prep.

• THE MORNING OF THE PROCEDURE

Beginning 6 hours prior to your **scheduled procedure time: (second dose)**

1. Take the Ondansetron (anti-nausea) 30 minutes prior to taking Dose 2 of the Suflave (bowel prep).
2. Fill second provided bottle with either provided flavor packet (lemon lime flavor) or any water enhancers of your choice like Crystal Light, MiO or Water Drops. Add lukewarm water to the top line of the container (at least 32 ounces), mix with spoon or shake until completely dissolved. You can mix a few hours prior to drinking and refrigerate to improve taste.
3. Drink **ALL** the solution, 8 ounces every 15 minutes. This can be done gradually to decrease the risk of nausea.
4. Drink an additional 16 oz or more of clear liquids of your choice. You may have clear liquids up to **4 hours** of your **scheduled procedure time**, then **nothing by mouth after that**.
5. **DO NOT TAKE** the following: Amaryl (glimepiride), Glucotrol (glipizide), Metformin (glucophage).
6. **DO NOT TAKE Ozempic, Mounjaro, Wegovy, Trulicity, Semaglutide, Saxenda, Zepbound, Vitoza, Rybelus, Adlyxin, Byureon, and Byetta (morning of procedure). Can resume after procedure.**
7. You should take the necessary medications you normally take on the morning of the procedure with **SIPS** of water to swallow medications.
8. Arrive at your scheduled **arrival time**.
9. Bring a list of your medications, insurance card and photo ID.
10. You must have a responsible adult drive you home after your procedure due to the sedation. It is best if your driver waits for you at the endoscopy center. **Taking a Cab, Uber, Lyft or Metro Call A Ride is NOT an option.** If you do not have a driver, we will not be able to sedate you for your procedure and the procedure may be canceled.
11. You may not drive or work until the following day.

• IMPORTANT INFORMATION ABOUT SEDATION

Sedatives used for endoscopy include Propofol. These medications provide amnesia and pain relief. You are breathing on your own. You may be groggy after your procedure and you **CAN NOT** work or drive the rest of the day, a responsible adult must drive you home after your procedure.

• QUESTIONS OR PROBLEMS

1. Suflave hotline is (855) MYSUFLAVE or (855) 697-5368.
2. Daytime phone number (314) 997-0554
3. After hours number (314) 388-6578
4. Website address is www.gidoctor.net
5. Billing Contacts:
 - Specialists in Gastroenterology: Physician, pathology and anesthesia (833) 415-5585
 - Advanced Endoscopy Center (314) 400-9999
 - Missouri Baptist Medical Center (314) 747-8845, will need procedure code (CPT code)